

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
18 March 2004 (18.03.2004)

PCT

(10) International Publication Number
WO 2004/022698 A2

(51) International Patent Classification⁷:

C12N

(74) Agent: BEN-AMI, Paulina; Ben-Ami & Associates, P. O.
BOX 94, 76100 Rehovot (IL).

(21) International Application Number:

PCT/IL2003/000725

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(22) International Filing Date:

3 September 2003 (03.09.2003)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

151660

9 September 2002 (09.09.2002) IL

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (*for all designated States except US*): BEN GURION UNIVERSITY OF THE NEGEV RESEARCH AND DEVELOPMENT AUTHORITY [IL/IL]; P.O. Box 653, 84105 Beer-Sheva (IL).

Published:

— *without international search report and to be republished upon receipt of that report*


(72) Inventors; and

(75) Inventors/Applicants (*for US only*): KUSHMARO, Ariel [IL/IL]; 19 Dekel St., 72905 Moshav Nir-Zvi (IL). GERESH, Shimonon [IL/IL]; 27 Tapuz St., 84965 Omer (IL).

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: METHOD FOR ISOLATING AND CULTURING UNCULTURABLE MICROORGANISMS

Polymeric membrane



Agar sphere

(57) Abstract: The invention provides a method for isolating and culturing a previously unculturable microorganism, which comprises: (i) collecting a sample from an environmental source; (ii) counting/estimating the number of microorganisms in the sample; (iii) diluting the sample in an appropriate medium; (iv) adding a gelating agent such as to entrap one or more microorganisms within a sphere of the gelating agent; (v) coating the spheres containing the entrapped microorganism(s) with a natural or synthetic polymer to form a polymeric membrane; (vi) incubating the coated spheres in the original environment for an appropriate time; (vii) cutting the spheres and scanning for microorganisms colonies; and (viii) isolating the microorganisms, and repeating steps (iii) to (vii) until a pure clone of said previously unculturable microorganism is obtained.

BEST AVAILABLE COPY

WO 2004/022698 A2

METHOD FOR ISOLATING AND CULTURING UNCULTURABLE MICROORGANISMS

5 FIELD OF THE INVENTION

The present invention relates to a method for isolating and culturing novel "uncultivable" microorganisms.

BACKGROUND OF THE INVENTION

10 Cultured microorganisms are the most common source of antibiotics and other medicinal agents. However, only a small percentage (less than 1%) of the viable bacteria in soil can be cultured on known nutrient media using current techniques such as petri dishes (Handelsman et al. 1998; Amann et al. 1995). The other 99% of uncultured/uncultivable microorganisms, with their genetic and
15 biochemical diversity, may emerge as a major source of new natural chemical structures that may be useful for humans, for example as drugs.

The exploration of previously uncultured microorganisms for the discovery of new useful natural products is now being carried out in several laboratories. The main approach involves genomics techniques such as the approach designated
20 metagenomics for the analysis of the collective genomes of the microorganisms in the soil community. According to this approach, DNA in large segments is cloned directly from soil into a culturable host and a sequence-based and functional genomic analysis is conducted on it. The intention is the isolation of new signals, new secondary metabolites that might have utility for humans, and the
25 reconstruction of an entire genome of an uncultured organism.

Molecular microbial ecology represents a recent development in research methods. It consists of utilizing techniques of molecular biology to investigate the ecology of microorganisms, and offers new tools to facilitate the detection and identification of microorganisms in the environment.

Molecular microbial ecology allowed the development of tools to address a central dogma of microbial ecology: an inability to cultivate more than a small proportion (0.1–1%) of the bacteria that can be visualized by direct count procedures (Head et al., 1998). Thus, the identification of bacteria by molecular methods represents an indispensable addition to the traditional methods based on the analyses of morphological and physiological characteristics. Among these culture-independent new techniques, the technique based on direct sequencing seems to be the most effective. It consists in sequencing a specific region of the bacterial chromosome, namely the bacterial 16S rDNA region, and in comparing this sequence with known sequences stored in data banks. All microorganisms possess at least one copy of the genes coding for the ribosomal RNA (rRNA), which are indispensable in any cells for the biosynthesis of proteins. Within these genes, the 16S rDNA region is principally used for the determination of the genus and the species of bacteria. By using this approach, it could be determined in many environmental samples the predominance of many different uncultured species. It might be feasible that the yet uncultured types of bacteria might be grown under laboratory conditions if just the right nutrients are found (Amann et al. 1995; Felske et al. 1999).

Recently, Kaeberlein et al. (2002, and published US Patent Applications Nos. 2003/0059866 and 2003/0059867) disclosed a new method for isolating and growing uncultivable microorganisms in pure culture in a simulated natural environment using a diffusion chamber. Microorganisms were separated from intertidal marine sediment particles, serially diluted, mixed with warm agar made with seawater, and placed in the diffusion chamber. The membranes allowed exchange of chemicals between the chamber and the environment, but restricted movement of cells. After the first membrane was affixed to the base of the chamber, the agar with microorganisms was poured in, and the top was sealed with another membrane (See Fig. 1, Kaeberlein et al., 2002' and Fig. 1a, US Patent Application No. 2003/0059866). The diffusion chamber consists of a stainless steel washer (70 mm o.d., 33 mm i.d., 3 mm in thickness) sandwiched between two 0.03- μ m pore-

size polycarbonate membranes. The membranes were glued to the washer forming the inner space filled with test microorganisms in semi-solid agar. The sealed chambers were placed on the surface of the sediment collected from the tidal flat and kept in a marine aquarium. Colonies of representative marine microorganisms were isolated in pure culture. These isolates did not grow on artificial media alone but formed colonies in the presence of other microorganisms.

Zengler et al. (2002) disclose a universal method that provides access to the immense reservoir of untapped microbial diversity. They utilized a technique that combines encapsulation of cells in gel microdroplets (using the OneCell System technology) for massively parallel microbial cultivation under low nutrient flux conditions, followed by flow cytometry to detect microdroplets containing microbial microcolonies.

In summary, most microorganisms in the environment have been overlooked as yet due to their resistance to cultivation on artificial media. The cultured microorganisms represent only a small fraction of natural microbial communities and hence the microbial diversity in terms of species richness and species abundance is grossly underestimated. Our understanding of microbial diversity is not represented by the cultured fraction of the microbial community (Wintzingerode et al., 1997).

SUMMARY OF THE INVENTION

It is an object of the present invention to provide a method for isolating and culturing previously unculturable microorganism.

In one preferred embodiment of the invention, the sample is collected from an environmental source and diluted in an appropriate medium after counting/estimating the number of microorganisms in the sample. A gelating agent is then added such as to entrap one or more microorganisms within a sphere of the gelating agent. The spheres containing the entrapped microorganism(s) are then coated with a natural or synthetic polymer to form a polymeric membrane. The

coated spheres with the entrapped microorganisms are incubated in the original environment and, after an appropriate time, are cut and scanned for microorganisms colonies. The microorganisms are isolated and subjected one or more times to the steps of dilution in an appropriate medium, entrapping in a gelating agent, coating
5 of the spheres containing the entrapped microorganisms, incubation of the coated spheres, cutting the spheres and scanning for microorganisms colonies, until a pure clone of said previously unculturable microorganism is obtained.

The present invention further provides a library of microorganisms obtained by the method of the invention, wherein each of the microorganisms was isolated
10 from a sphere as described.

The present invention still further provides a method for screening and identification of new drugs and other substances of commercial interest in the pharmaceutical, chemical, biotechnology and other industries as well as in the agriculture, which comprises cultivating a previously unculturable microorganism
15 or screening a library of previously unculturable microorganisms, and isolating and identifying compounds having biological or other activity of interest.

BRIEF DESCRIPTION OF THE DRAWINGS

Fig. 1 depicts a schematic drawing of an agar sphere coated with a polymeric
20 membrane.

Fig. 2 is a scanning electron microscopy photograph showing an agar sphere coated with a polysulfone membrane as obtained in Example 1 hereinafter.

Fig. 3 is a photograph (x1000) of microcolony of an unidentified rod-shaped bacterium obtained from soil as described in Example 3 hereinafter.

DETAILED DESCRIPTION OF THE INVENTION

The present invention relates to a method for isolation and culture of microorganisms suitable for microorganisms from any environmental source.

According to one aspect, the present invention relates to a method for isolating and culturing a previously unculturable microorganism, which comprises:

- (i) collecting a sample from an environmental source;
- (ii) counting/estimating the number of microorganisms in the sample;
- (iii) diluting the sample in an appropriate medium;
- (iv) adding a gelating agent such as to entrap one or more microorganisms within a sphere of the gelating agent;
- (v) coating the spheres containing the entrapped microorganism(s) with a natural or synthetic polymer to form a polymeric membrane;
- (vi) incubating the coated spheres in the original environment for an appropriate time;
- (vii) cutting the spheres and scanning for microorganisms colonies; and
- (viii) isolating the microorganisms, and repeating steps (iii) to (vii) until a pure clone of said previously unculturable microorganism is obtained.

According to the invention, an environmental sample is collected and the number of bacteria is counted/estimated by microscopic observation, namely, all cells are counted in a large square: 12 cells (in practice, several squares are counted and the numbers averaged). The sample is diluted 1/10 in an appropriate medium, 1 ml of the broth is transferred to a tube, diluted again to 1/10², 1 ml thereof is transferred to another tube, and the dilution steps are repeated until about 1/10⁶, when approximately one bacterium will be entrapped in the gelating agent sphere. The gelating agent sphere is coated with a polymeric membrane and incubated in the environment for weeks to months. At the end of the incubation period, the coated sphere is cut, the bacterium is then isolated, cultured in the sphere, and subjected to molecular biology techniques.

The environmental source from which the samples are collected may be any

terrestrial, aquatic or marine source such as soil, biofilms, sediments (e.g. coral or other marine sediments, aquifer sediments and the like), waste waters and the like. The sample is suspended in its natural or other appropriate medium, and is divided, for example, into 1-ml tubes, and each divisional sample is then subjected to
5 counting/estimation of the number of microorganisms by well-known techniques, for example by DAPI (4',6-diamidino-2-phenylindole) staining of the cells and direct microscopic count of the DAPI-stained cells.

In the next step, the samples are diluted as necessary in an appropriate medium. As used herein, an "appropriate medium" is intended to mean a medium
10 compatible with the environment from which the sample has been collected with respect to physico-chemical parameters such as pH, salinity, temperature, oxygen concentration, and the like. The medium may be sterile water, sterile saline, sterile water containing suitable ingredients for compatibility with the environmental source, and the like. For example, a sample collected from a marine source will
15 have the salinity corresponding to the marine source and the salt concentration will be higher if the sample is originated from the Dead Sea. When the sample is collected from soil, the medium may be sterile water.

The next step consists in the addition of a gelating agent as a matrix to the diluted samples. Any suitable natural, semi-synthetic or synthetic gelating agent
20 may be used such as, but not limited to, agar, alginate, carrageenans, gum Arabic, guar gum, traganth gum, xanthan gum, propyleneglycolalginate, and microcrystalline cellulose.

In one preferred embodiment, the gelating agent is agar. Preferably, the diluted samples are mixed with warm (40-50°C) autoclaved agar (0.7-2%) such as
25 to entrap preferably one, or more, microorganisms within a sphere of the gelating agent. The agar spheres can be obtained by dripping agar droplets into cold mineral oil. The sphere size of 0.1 mm or less, up to 5 mm, preferably up to 2-3 mm, in diameter, can be determined by the nozzle diameter and dripping rate.

The gelating agent spheres containing the entrapped microorganism(s) are
30 then coated with a natural or synthetic polymer such as, but not limited to, a

polysulfone, an alginate, an epoxy resin, polyacrylamide, silica gel and the like, such as to form a multilayered membrane. Polysulfone resin of average m.w. 35000, Na-alginate, and epoxy resin such as Epikote 255 have been disclosed before for biomass entrapment for different applications (Ferguson et al., 1989; Blanco et al., 1999) and can be used according to the invention.

For coating, the polymer is first dissolved in a suitable solvent, the dried gelating agent spheres containing the entrapped microorganism are introduced into the polymer solution, and are then transferred into a medium that enables coating of the spheres by several layers of the polymer, thus forming the desired spheres coated by the polymeric membrane. In one embodiment, agar spheres containing one or more microorganisms are immersed into a solution of polysulfone in dimethylformamide (DMF) and transferred to water in order to obtain the desired polymeric coating. The polymeric membrane allows exchange of chemicals between the sphere and the environment, but restrict movement of cells. The polymer can be opaque, but preferably it is transparent such that the colonies inside the sphere can be seen.

The next step consists in the incubation of the polymeric coated gelating agent spheres containing one or more microorganisms in the environment from which the original sample has been collected, for an appropriate time. This is the alternative to cultivate such microorganisms that cannot grow in known growth media for microorganisms. The incubation in the environment can take from days to months. When the sample is collected from a marine source, the incubation may be carried out in a container, e.g., aquarium, containing water with the degree of salinity of the marine source. When the sample is collected from soil, the incubation may be carried out in containers, e.g. pots, filled with the same soil material.

After the incubation, the spheres are cut and placed on a glass slide and covered with a coverslip, and the entire volume of agar is scanned for microbial colonies at magnification of 400X and 1,000X. The microorganisms are then isolated into pure culture by successive re-inoculation of individual colonies into new spheres followed by incubation in the environment, by repetition of steps (iii)

to (vii) as many times as necessary, until a pure clone of the previously unculturable microorganism is obtained.

After incubation in the environment, previously uncultured microorganisms can be isolated and then subjected to molecular biology and genomics techniques, and/or cultured for the production of bioactive materials. Libraries can be construed
5 composed of microorganisms, each isolated from a separate sphere, and can be used for identification of new biologically active compounds, even without identification and characterization of the microorganisms. When the identified biologically active compound is a small organic molecule, its structure can be determined by known
10 methods, it is then synthesized, the biological activity is ascertained and it can then be formulated in pharmaceutical or veterinary compositions.

The method of the present invention allows isolation/identification of new types of microorganisms, such as bacteria, previously considered as unculturable, and the establishment of libraries of "uncultivable" microorganisms useful for the
15 drug discovery and biotechnology industries. The method enables exploration of new natural products from previously uncultured microorganisms. New genes might be obtained from the previously uncultivable microbial communities, and new biologically active materials such as proteins, enzymes and antibiotics of utility to humans may be discovered.

20 According to the invention, metagenomics techniques can be used to address the genetic structure and functional composition of a sample irrespective of whether the microorganism can be cultured. Molecular methodologies such as PCR of select molecular targets can be used to discover genes with useful properties. Microbial communities can be profiled by techniques well known in the art. Cloning and
25 sequencing of molecular targets such as 16S rDNA enable identification of indigenous and novel organisms.

In one embodiment of the invention, cells or extracts from unculturable microorganisms are subjected to analysis by 16S RNA gene sequencing. Ribosomal RNA genes from the samples, microcolonies or cultures are amplified by PCR by
30 using specific 16S RNA oligonucleotide primers for bacteria. After cloning the PCR

products, the inserts are screened by their restriction patterns (RFLP - restriction fragment length polymorphism). The clones are submitted to sequence analysis and compared with known 16S RNA genes using, for example, the online GenBank database (<http://ncbi.nlm.nih.gov/GenBank>). In this way, it can be determined whether or not the microorganism represents a new species/genus.

5 The present invention further provides a method for genomic characterization of previously unculturable microorganisms, which comprises: (i) collecting a sample from an environmental source; (ii) counting/estimating the number of microorganisms in the sample; (iii) diluting the sample in an appropriate medium; (iv) adding a gelating agent such as to entrap one or more microorganisms within a sphere of the gelating agent; (v) coating the spheres containing the entrapped microorganism(s) with a natural or synthetic polymer to form a polymeric membrane; (vi) incubating the coated spheres in the original environment for an appropriate time; (vii) cutting the spheres and extracting the microorganisms by chemical lysis using an agent for extraction of genomic DNA; (viii) processing the total genomic DNA to establish the restriction fragment length polymorphism (RFLP) pattern of the microorganisms; (ix) analyzing the RFLP patterns to identify unique clones that are submitted to sequence analysis; and (x) identifying the microorganisms by comparison of these sequences with sequences available at the GenBank database.

20 In one preferred embodiment, a method is provided wherein the microorganisms are isolated from a marine source, which comprises: (i) collecting a sample from a marine source; (ii) counting/estimating the number of microorganisms in the sample; (iii) diluting the sample in sterile seawater; (iv) adding a gelating agent such as to entrap one or more microorganisms within a sphere of the gelating agent; (v) coating the spheres containing the entrapped microorganism(s) with a natural or synthetic polymer to form a polymeric membrane; (vi) incubating the coated spheres in an aquarium containing seawater for an appropriate time; (vii) cutting the spheres and extracting the microorganisms by chemical lysis using an agent for extraction of genomic DNA; (viii) processing

25
30

the total genomic DNA to establish the restriction fragment length polymorphism (RFLP) pattern of the microorganisms; (ix) analyzing the RFLP patterns to identify unique clones that are submitted to sequence analysis; and (x) identifying the microorganisms by comparison of these sequences with sequences available at the GenBank database.

In a preferred embodiment of the present invention, the microorganism is a bacterium isolated from a marine source consisting of coral mucus. According to this embodiment, extracts from unculturable microorganisms from Red Sea coral mucus were amplified by PCR using the specific 16S RNA oligonucleotide primers for bacteria of SEQ ID NO:1 (forward primer, 8F) and SEQ ID NO:2 (reverse primer, 1512 R).

The most abundant sequence (70%) in this Red Sea coral mucus sample corresponded to the partial 16S rDNA sequence characterized by the SEQ ID NO: 3 (1,428 bp), and represent the Bacteria (domain), Proteobacteria (phylum), Betaproteobacteria (class), Burkholderiales (order), Alcaligenaceae (family), Alcaligenes (genus).

The next most abundant clone (20%) in the Red Sea coral mucus sample corresponded to the partial 16S rDNA sequence characterized by the SEQ ID NO:4 (1,382 bp), and represent the Bacteria (domain), Proteobacteria (phylum), Alphaproteobacteria (class), Rhodobacteriales (order), Rhodobacteraceae (family), Roseobacter (genus) or Ruegeria (genus) or unclassified (genus).

The less abundant clone (10%) in the Red Sea coral mucus sample corresponded to the partial 16S rDNA sequence characterized by the SEQ ID NO: 5 (1,483 bp), and represent the Bacteria (domain), Proteobacteria (phylum), Gammaproteobacteria (class), Endobacteriales (order) or Verrucomicrobiales (order).

In another embodiment of the present invention, the microorganism is isolated from a soil sample. According to this embodiment, a clone from a library constructed from unculturable soil bacteria from the soil of the Halutza region, Negev, Israel, was amplified by PCR using the specific 16S RNA oligonucleotide

primers for bacteria of SEQ ID NO:1 (forward primer, 8F) and SEQ ID NO:2 (reverse primer, 1512 R).

The most abundant clone in the Halutza soil bacteria corresponded to the partial 16S rDNA sequences characterized by the SEQ ID NO: 6 (730 bases, by direct 8F primer) and SEQ ID NO: 7 (689 bases, by reverse 1512R primer) and represent the Proteobacteria (phylum), Betaproteobacteria (class), Burkholderiales (order), Alcaligenaceae (family), Alcaligenes (genus).

The next most abundant clone in this sample of Halutza soil bacteria corresponded to the partial 16S rDNA sequences characterized by the SEQ ID NO:8 (651 bases, by direct 8F primer) and SEQ ID NO: 9 (886 bases, by reverse 1512R primer) and represent the Proteobacteria (phylum), Gammaproteobacteria (class), Pseudomonadales (order), Pseudomonadaceae (family), Pseudomonas (genus).

In another aspect, the present invention relates to a previously uncultured microorganism isolated by the method of the present invention. The microorganism may be a eukaryote, e.g. a fungus, or a prokaryote, e.g., a bacterium. In a preferred embodiment, the microorganism is a bacterium. In most preferred embodiments, the bacterium is the bacterium isolated from soil depicted in Fig. 3, or it is a bacterium comprising the partial 16S rDNA nucleic acid sequences selected from the group consisting of SEQ ID NO:3, SEQ ID NO:4, and SEQ ID NO:5, or the pair of 16S rDNA sequences consisting of SEQ ID NO:6 and NO:7, or SEQ ID NO:8 and NO:9.

In a further aspect, the present invention relates to a library of previously uncultured microorganisms obtained by the method of the invention and to the use of said library for the discovery of new biologically active agents including, but not being limited, to new antibiotics, enzymes, biocatalysts, genes.

It is further envisaged to construct BAC, cosmid and small insert libraries from diverse environmental samples and then subject the libraries to a screening for novel genes, proteins and small molecules exhibiting activities of interest. For example, 16S rRNA gene clone libraries can be formed from mixed colonies of microorganisms and screened.

The colonies of microorganisms can also be screened for antibiotic activity by contacting diluted samples with a strain of interest and studying the influence on its growth. Colonies of unculturable microorganisms that produce compounds with antibiotic activity will inhibit growth of strains. Said compounds can then be isolated, purified, analyzed and either synthesized for use as antibiotic or used as a model for further drug discovery.

Libraries of test extracts of the microorganisms can also be tested for activity by automated highthroughput biochemical or biological assays using, for example, a panel of test microorganisms to test antibiotic activity, or a panel of enzymes or antibodies to find compounds that affect their activities.

The invention will now be illustrated by the following non-limiting Examples.

EXAMPLES

Example 1: Laboratory scale wastewater bioreactor

An environmental sample was obtained from laboratory scale wastewater bioreactor (waste water from Ramat Hovav Toxic Waste Dumping Site, Negev, Israel) and estimated for microorganism number by DAPI-staining and microscope direct counting (10^8 - 10^9 cell/mL). The sample was diluted 8- and 9-fold with water in order to entrap approximately one microorganism in one agar sphere.

The diluted samples were mixed with warm (50°C) autoclaved agar (DIFCO) (900 µl agar per 100 µl diluted sample, final concentration 1.5 % agar). Agar spheres of approximately 1-2 mm in diameter containing the entrapped microorganism(s) were formed by dripping droplets of the mixture into cold mineral oil.

A solution of 10% polysulfone of m.w. ca. 35,000 (Sigma-Aldrich, Product No. 42,830-2) in DMF was prepared and used to coat the dried agar spheres containing the entrapped microorganism(s). For this purpose, the agar spheres were introduced into the polymer solution, and then transferred into water in order to

obtain the desired polymeric membrane. Fig. 2 is a photograph of such a coated agar sphere.

5 The polysulfone-coated agar spheres containing the entrapped microorganism(s) were then incubated for 3 weeks in a wastewater bioreactor. At the end of the incubation, the agar spheres were cut and placed between a glass slide and a coverslip, and the entire volume of agar was scanned for microbial colonies at magnification of 400X and 1000X.

10 In the spheres, we could see development of isolated colonies of bacteria and fungi presumably uncultivable, since experiments to grow them in ordinary agar plates were not successful and no bacterial growth was observed.

Example 2: Microbial communities inhabiting coral mucus

15 Mucus of coral heads from the Red Sea was sampled by collecting ca. 1 ml¹ in a sterile disposable 50-ml polypropylene centrifuge tube. Mucus bacteria were counted directly under phase microscopy. Mucus was diluted by sterile seawater and placed on marine agar for culturable colony forming units (CFU). Several dilutions (10^{-7} , 10^{-6} , 10^{-5} and 10^{-4}) were mixed with 2% marine agar in ratio 1 to 1 and agar spheres were coated with polysulphone and incubated in a seawater aquarium containing corals for several weeks. Agar spheres of approximately 1-2
20 mm in diameter containing the entrapped microorganism(s) were formed by dripping droplets of the mixture into cold mineral oil.

A solution of 10% polysulfone of m.w. ca. 35,000 (Sigma-Aldrich, Product No. 42,830-2) in DMF was prepared and used to coat the dried agar spheres containing the entrapped microorganism(s). For this purpose, the agar spheres were
25 introduced into the polymer solution, and then transferred into water in order to obtain the desired polymeric membrane.

The polysulfone-coated agar spheres containing the entrapped microorganism(s) were then incubated for 3 weeks in a seawater aquarium containing corals. At the end of the incubation, the agar spheres were cut and placed
30 between a glass slide and a coverslip, and the entire volume of agar was scanned for

microbial colonies at magnification of 400X and 1000X.

In the spheres, we could see development of isolated colonies of bacteria presumably uncultivable, since experiments to grow them in ordinary agar plates were not successful and no bacterial growth was observed.

5 Agar spheres including unculturable bacteria were extracted by chemical lysis protocol using CTAB solution (hexadecyltrimethylammonium bromide, Sigma H-5882) for genomic DNA. The total DNA was amplified with a Mastercycler gradient thermocycler (Eppendorf, Westbury, N.Y.) by PCR using specific 16S rRNA primers for bacteria [forward primer, 8F (5'-GGATCCAGACTTT
10 GAT(C/T)(A/C)TGGCTCAG – SEQ ID NO:1) and reverse primer, 1512R (5' GTGAAGCTTA CGG(C/T)TAGCTTGTTACGACTT – SEQ ID NO:2)]. Primers used in the PCR amplifications were obtained from Sigma-Genosys. Reaction mixtures included a 12.5 µl ReddyMix (PCR Master mix containing 1.5 mM MgCl₂ and 0.2 mM concentration of each deoxynucleoside triphosphate) (ABgene, Surrey,
15 UK), a 1 pmol each of the forward and reverse primers, 1 µl of the sample preparation, and water to bring the total volume to 25 µl. An initial denaturation-hot start of 4 min at 95°C was followed by 30 cycles of the following incubation pattern: 94°C for 20 sec, 56°C for 20 sec, and 72°C for 105 sec. A final soak at 72°C for 20 min concluded the reaction.

20 PCR products were purified by electrophoresis through a 0.8% agarose gel (Sigma), stained with ethidium bromide, and visualized on a UV transilluminator. The approximately 1,500-bp heterologous ribosomal DNA (rDNA) product was excised from the gel, and the DNA was purified from the gel slice by using the QIAquick gel extraction kit (Qiagen, Germany). The gel-purified PCR product was
25 cloned into the pDrive vector by QIAgen PCR cloning kit (Qiagen, Germany) and transformed into calcium chloride-competent XL MRF' *E.coli* cells. Plasmid DNA was isolated from individual clones by QIAprep Spin Miniprep kit (Qiagen, Germany).

Aliquots from a subset of the samples of purified plasmid DNA were digested
30 with 5 U of the restriction enzyme *EcoRI* for more than 4 h at 37°C, and the

digested product was separated by electrophoresis on a 1% agarose gel. After staining with ethidium bromide, the bands were visualized on a UV transilluminator and the RFLP (restriction fragment length polymorphism) patterns were analyzed to select clones containing the appropriately sized insert. Plasmid DNA from these clones was then digested with the six-base recognition site enzymes *Sph*I, *Sal*I, *Eco*RV, *Bam*HI, *Hind*III, *Not*I, *Xba*I and *Xho*I (Sigma) under the conditions described above. The digest products were then separated by electrophoresis on a 1% agarose gel, stained with ethidium bromide, and the RFLP patterns were used to identify unique clones to be submitted for sequence analysis.

Three additional internal primers were designed for completed sequencing of the clones. Sequence reactions were performed on the plasmid templates by using an ABI 3700 capillary sequencer. The rDNA sequences were first compared with those in the GenBank database with the basic local alignment search tool (BLAST) network service through San-Diego Supercomputer Center (SDSC). From the alignments created by this search, the orientation of each cloned 16S rRNA gene could be determined and a rough phylogenetic association was established. Each sequence was analyzed using the CHIMERA CHECK program (version 2.7) available at the Ribosomal Database Project.

Clone library constructed from agar spheres (10 colonies) inoculated by mucus, revealed three different patterns by RFLP by ratio 7 to 2 to 1. Representative DNA from each group was partial sequenced by using direct (8F) and reversed (1512R) primers. Microorganism identification was based on comparison of these sequences with the GenBank database and exhibited high division-level diversity of bacterial sequences.

The most abundant clone (70%) is characterized by partial 16S rDNA sequence of SEQ ID NO:3 that represents Bacteria (domain), Proteobacteria (phylum), Betaproteobacteria (class), Burkholderiales (order), Alcaligenaceae (family), Alcaligenes (genus). The next most abundant clone (20%) is characterized by partial 16S rDNA sequence of SEQ ID NO:4 that represents Bacteria (domain), Proteobacteria (phylum), Alphaproteobacteria (class), Rhodobacteriales (order),

Rhodobacteraceae (family), Roseobacter (genus) or Ruegeria (genus) or unclassified (genus). The less abundant clone (10%) in the Red Sea coral mucus sample is characterized by partial 16S rDNA sequence of SEQ ID NO:5 that represents Bacteria (domain), Proteobacteria (phylum), Gammaproteobacteria (class), Endobacteriales (order) or Verrucomicrobiales (order).

Example 3: Microbial communities inhabiting soil

A sample was collected from the soil of the Halutza region, Negev, Israel, and processed as described in Example 2. The medium was sterile tapwater and the incubation of the polysulfone-coated agar spheres was carried out in pots filled with soil material from the Halutza region. A clone library constructed from agar spheres (10 colonies) inoculated with the soil bacteria was incubated for 3 weeks, analyzed as described above and revealed two different patterns by RFLP by ratio 9 to 1. DNA from two clones were partially sequenced by using direct (8F) and reversed (1512R) primers and microorganism identification was based on comparison of these sequences with the GenBank database.

The most abundant sequence (90%) of this sample of soil bacteria is characterized by partial 16S rDNA sequences of SEQ ID NO:9 and SEQ ID NO:10, that represent Proteobacteria (phylum), Betaproteobacteria (class), Burkholderiales (order), Alcaligenaceae (family), Alcaligenes (genus). The less abundant (10%) clone is characterized by partial 16S rDNA sequences of SEQ ID NO:11 and SEQ ID NO:12, that represent Proteobacteria (phylum), Gammaproteobacteria (class), Pseudomonadales (order), Pseudomonadaceae (family), Pseudomonas (genus).

Fig. 3 is a photograph (x1000) of a microcolony of an unidentified rod-shaped bacterium obtained in purified form from the Halutza soil.

REFERENCES

Amann, R.I., Ludwig, W. and Schleifer, K-H. (1995) Phylogenetic identification and in situ detection of individual microbial cells without cultivation. *Microbiol. Rev.* 59, 143-169.

5 Blanco A., Sanz B., Llama J. M., Serra L. M.. (1999). Biosorption of heavy metals to immobilised *Phormidium laminosum* biomass. *Journal of Biotechnology* 69: 227-240.

 Felske A., Wolterink A., van Lis R., W. M. de Vos, A. D.L. Akkermans. (2000). Searching for predominant soil bacteria: 16S rDNA cloning versus strain
10 cultivation. *FEMS Microbiology Ecology* 30:137-145.

 Ferguson C.R., Peterson M.R., Jeffers T.H. (1989). Removal of metal contaminants from waste waters using biomass immobilized in polysulfone beads. In: Scheiner, B.J., Doyle, F.M., Kawatras, S.K. (Eds.), *Biotechnology in Minerals and Metal Processing*. Society of Mining Engineers, Littleton, CO, pp. 193-199.
15

 Handelsman J., Rondo R. M., Brady F. S., Clardy J., Goodman M. R. (1998). Molecular biological access to the chemistry of unknown soil microbes: a new frontier for natural products. *Chem. Biol.* 5(10):245-249.

20 Head I.M., Saunders J.R., Pickup R.W. (1998). Microbial Evolution, Diversity, and Ecology: A Decade of Ribosomal RNA Analysis of Uncultivated Microorganisms. *Microb Ecol* 35:1-21.

 Kaeberlein T., Lewis K., Epstein S. (2002). Isolating "uncultivable"
25 microorganisms in pure culture in a simulated natural environment. *Science* 296: 1127-1129.

 Spring S., Schulze R., Overmann J., Schleifer K. (2000). Identification and

characterization of ecologically significant prokaryotes in the sediment of freshwater lakes: molecular and cultivation studies. FEMS Microbiology Reviews 24:573-590.

- 5 Wintzingerode F., Goebel U., Stackebrandt E. (1997). Determination of microbial diversity in environmental samples: pitfalls of PCR-based rRNA analysis. FEMS Microbiology Reviews 21:213-229.
- Zengler K., Toledo G., Rappe M., Elkins J., Mathur J. E., Short J.M., and
10 Keller M. (2002). Cultivating the uncultured. PNAS 99 (24): 15681-15686.

15

20

CLAIMS:

1. A method for isolating and culturing a previously unculturable microorganism, which comprises:

- 5 (i) collecting a sample from an environmental source;
- (ii) counting/estimating the number of microorganisms in the sample;
- (iii) diluting the sample in an appropriate medium;
- 10 (iv) adding a gelating agent such as to entrap one or more microorganisms within a sphere of the gelating agent;
- (v) coating the spheres containing the entrapped microorganism(s) with a natural or synthetic polymer to form a polymeric membrane;
- 15 (vi) incubating the coated spheres in the original environment for an appropriate time;
- (vii) cutting the spheres and scanning for microorganisms colonies; and
- 20 (viii) isolating the microorganisms, and repeating steps (iii) to (vii) until a pure clone of said previously unculturable microorganism is obtained.

2. A method according to claim 1 wherein said environmental source is a terrestrial, aquatic or marine source.

25 3. A method according to claim 1 or 2 wherein said appropriate medium of (iii) is a medium compatible with the environment from which the sample has been collected.

4. A method according to claim 1 wherein said gelating agent is a natural, semi-synthetic or synthetic gelating agent selected from agar, alginate, carrageenans, gum Arabic, guar gum, traganth gum, xanthan gum, propyleneglycolalginate, and mycrocrystalline cellulose, and is preferably agar.

5

5. A method according to claim 1 wherein said gelating agent sphere of (iv) has a size from 0.1 mm or less to about 5 mm, preferably 1-2 mm in diameter.

6. A method according to claim 1 wherein said natural or synthetic polymer for
10 coating the spheres containing the microorganism(s) is a natural or synthetic transparent or opaque polymer selected from a polysulfone, an alginate, an epoxy resin, a polyacrylamide, silica gel polysulfone, alginate and epoxy resin.

7. A microorganism isolated by the method according to claim 1.

15

8. A microorganism according to claim 7 consisting of a bacterium.

9. A library of microorganisms according to claim 7 or 8.

10. A method for genomic characterization of previously unculturable
20 microorganisms, which comprises: (i) collecting a sample from an environmental source; (ii) counting/estimating the number of microorganisms in the sample; (iii) diluting the sample in an appropriate medium; (iv) adding a gelating agent such as to entrap one or more microorganisms within a sphere of the gelating agent; (v) coating the spheres containing the entrapped microorganism(s) with a natural or
25 synthetic polymer to form a polymeric membrane; (vi) incubating the coated spheres in the original environment for an appropriate time; (vii) cutting the spheres and extracting the microorganisms by chemical lysis using an agent for extraction of genomic DNA; (viii) processing the total genomic DNA to establish the restriction fragment length polymorphism (RFLP) pattern of the microorganisms;
30 (ix) analyzing the RFLP patterns to identify unique clones that are submitted to

sequence analysis; and (x) identifying the microorganisms by comparison of these sequences with sequences available at the GenBank database.

11. A method according to claim 10 wherein the microorganisms are isolated from a marine source, which comprises: (i) collecting a sample from a marine source; (ii) counting/estimating the number of microorganisms in the sample; (iii) diluting the sample in sterile seawater; (iv) adding a gelating agent such as to entrap one or more microorganisms within a sphere of the gelating agent; (v) coating the spheres containing the entrapped microorganism(s) with a natural or synthetic polymer to form a polymeric membrane; (vi) incubating the coated spheres in an aquarium containing seawater for an appropriate time; (vii) cutting the spheres and extracting the microorganisms by chemical lysis using an agent for extraction of genomic DNA; (viii) processing the total genomic DNA to establish the restriction fragment length polymorphism (RFLP) pattern of the microorganisms; (ix) analyzing the RFLP patterns to identify unique clones that are submitted to sequence analysis; and (x) identifying the microorganisms by comparison of these sequences with sequences available at the GenBank database.

12. A bacterium isolated by the method according to claim 11.

13. A method according to claim 11 wherein the marine source is coral mucus.

14. A bacterium isolated from coral mucus by the method of claim 13 and characterized by partial 16S rDNA sequence selected from the group consisting of SEQ ID NO:3, SEQ ID NO:4, and SEQ ID NO:5.

15. A method according to claim 10 wherein the microorganisms are isolated from a soil source.

16. A bacterium isolated from soil by the method according to claim 15 and characterized by partial 16S rDNA sequences selected from the group consisting of

the pair of sequences of SEQ ID NO:6 and SEQ ID NO:7, and SEQ ID NO:8 and SEQ ID NO:9.

17. Use of a library of microorganisms obtained by the method according to claim 1 for identification of new drugs.

ABSTRACT

5 The invention provides a method for isolating and culturing a previously
unculturable microorganism, which comprises: (i) collecting a sample from an
environmental source; (ii) counting/estimating the number of microorganisms in the
sample; (iii) diluting the sample in an appropriate medium; (iv) adding a gelating
agent such as to entrap one or more microorganisms within a sphere of the gelating
10 agent; (v) coating the spheres containing the entrapped microorganism(s) with a
natural or synthetic polymer to form a polymeric membrane; (vi) incubating the
coated spheres in the original environment for an appropriate time; (vii) cutting the
spheres and scanning for microorganisms colonies; and (viii) isolating the
microorganisms, and repeating steps (iii) to (vii) until a pure clone of said
15 previously unculturable microorganism is obtained.

1/2

Polymeric membrane




Fig. 1




Fig. 2

2/2



Fig. 3

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
18 March 2004 (18.03.2004)

PCT

(10) International Publication Number
WO 2004/022698 A3

(51) International Patent Classification⁷: **C12N 1/20**,
9/80, C12P 21/04, C12Q 1/02

(21) International Application Number:
PCT/IL2003/000725

(22) International Filing Date:
3 September 2003 (03.09.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
151660 9 September 2002 (09.09.2002) IL

(71) Applicant (for all designated States except US): **BEN GU-
RION UNIVERSITY OF THE NEGEV RESEARCH
AND DEVELOPMENT AUTHORITY** [IL/IL]; P.O. Box
653, 84105 Beer-Sheva (IL).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **KUSHMARO**,
Ariel [IL/IL]; 19 Dekel St., 72905 Moshav Nir-Zvi (IL).
GERESH, Shimona [IL/IL]; 27 Tapuz St., 84965 Omer
(IL).

(74) Agent: **BEN-AMI, Paulina**; Ben-Ami & Associates, P. O.
BOX 94, 76100 Rehovot (IL).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC,
SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA,
UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,
ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO,
SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM,
GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments

(88) Date of publication of the international search report:
6 January 2005

*For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*

(54) Title: **METHOD FOR ISOLATING AND CULTURING UNCULTURABLE MICROORGANISMS**

(57) Abstract: The invention provides a method for isolating and culturing a previously unculturable microorganism, which comprises: (i) collecting a sample from an environmental source; (ii) counting/estimating the number of microorganisms in the sample; (iii) diluting the sample in an appropriate medium; (iv) adding a gelating agent such as to entrap one or more microorganisms within a sphere of the gelating agent; (v) coating the spheres containing the entrapped microorganism(s) with a natural or synthetic polymer to form a polymeric membrane; (vi) incubating the coated spheres in the original environment for an appropriate time; (vii) cutting the spheres and scanning for microorganisms colonies; and (viii) isolating the microorganisms, and repeating steps (iii) to (vii) until a pure clone of said previously unculturable microorganism is obtained.

WO 2004/022698 A3

INTERNATIONAL SEARCH REPORT

International application No.

PCT/IL03/00725

A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : C12N 1/20, 9/80; C12P 21/04; C12Q 1/02

US CL : 435/29, 71.2, 228, 252.2

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 435/29, 71.2, 228, 252.2

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched
REGISTRY

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
Please See Continuation Sheet

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 5,744,325 A (FUJISHIMA et al.) 28 April 1998 (28.04.1998), Entire document, especially abstarct, Clumn 2, Lines 10-65	1-12
Y	US 5,032,508 A (NAUGHTON et al.) 16 July 1991(16.07.1991), Entire document.	1-17

☐ Further documents are listed in the continuation of Box C.

☐ See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T"

later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X"

document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y"

document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&"

document member of the same patent family

Date of the actual completion of the international search

16 August 2004 (16.08.2004)

Date of mailing of the international search report

29 NOV 2004

Name and mailing address of the ISA/US

Mail Stop PCT, Attn: ISA/US
Commissioner for Patents
P.O. Box 1450
Alexandria, Virginia 22313-1450

Facsimile No. (703) 305-3230

Authorized officer

Dr. Kailash C. Srivastava

Telephone No. (571)-272-1600

Janice Ford
for

INTERNATIONAL SEARCH REPORT

PCT/IL03/00725

Continuation of B. FIELDS SEARCHED Item 3:

USPT, PGPB, EPAB, JPAB, DWPI, BIOSIS, BIOTECHABS, BIOTECNO, CABA, CANCERLIT, CAPLUS, CEABA-VTB, DISSABS, EMBASE, ESBIODASE, FROSTI, IFIPAT, HCAPLUS, IICST-PLUS, LIFESCI, MEDICINF, MEDLINE, PHIN, PROMPT, RDISCLOSURE, SCISEARCH, TOXCENTER, USPATFUL, WPIDS.

Search Strategy:

bacteria, microorganism, encapsulatd, gell, gum, synthetic or natural plymer, microcrystalline cellulose, isolate, terrestrial, environmental, aquatic, sediment, vibrio, medicinal, drug screening, cultur, gel medium

SEQUENCE LISTING

<110> BEN-GURION UNIVERSITY OF THE NEGEV RESEARCH AND DEVELOPMENT AUTHORITY
KUSHMARO, Ariel
GERESH, Shimona

<120> METHODS FOR ISOLATING AND CULTURING UNCULTURABLE MICROORGANISMS

<130> BGU-003 PCT

<150> IL 151660
<151> 2002-09-09

<160> 9

<170> PatentIn version 3.1

<210> 1
<211> 26
<212> DNA
<213> Synthetic Sequence

<220>
<221> misc_feature
<223> Residue no. 17 is C or T, residue no. 18 is A or C

<400> 1
ggatccagac tttgatcatg gctcag 26

<210> 2
<211> 30
<212> DNA
<213> Synthetic Sequence

<220>
<221> misc_feature
<223> Residue no. 14 is C or T

<400> 2
gtgaagctta cggctagctt gttacgactt 30

<210> 3
<211> 1428
<212> DNA
<213> Bacterium-partial 16S rDNA of unknown species from Red Sea coral mucus

<400> 3
agtcgaacgg cagcgcgaga gagcttgctt ttttggcggc gagtggcgga cgggtgagta 60
atatatcgga acgtgccag tagcggggga taactactcg aaagagtggc taataccgca 120
tacgccctac gggggaaagg gggggatcgc aagacctctc actattggag cggccgatat 180
cggattagct agttggtggg gttaaaggctc accaaggcaa cgatccgtag ctggtttgag 240
aggacgacca gccacactgg gactgagaca cggcccagac tcctacggga ggcagcagtg 300
gggaattttg gacaatgggg gaaaccctga tccagccatc ccgcgtgtat gatgaaggcc 360
ttcgggttgt aaagtacttt tggcagagaa gaaaaggtat cccctaatac gggatactgc 420
tgacggtatc tgcagaataa gcaccggcta actacgtgcc agcagccgcg gtaatacgta 480

gggtgcaagc gttaatcgga attactgggc gtaaagcgtg tgtaggcggt tcggaagaa 540
agatgtgaaa tcccaggggc tcaaccttgg aactgcattt ttaactgccg agctagagta 600
tgtcagaggg gggtagccat tccacgtgta gcagtgaat gcgtagatat gtggaggaat 660
accgatggcg aaggccgccc cctgggataa tactgacgt cagacacgaa agcgtgggga 720
gcaaacagga ttagataccc tggtagtcca cgccctaaac gatgtcaact agctgttggg 780
gccgttaggc cttagtagcg cagctaacgc gtgaagtga ccgcctgggg agtacggtcg 840
caagattaaa actcaaagga attgacgggg acccgacaaa gcggtggatg atgtggatta 900
attcgatgca acgcgaaaaa ccttacctac ccttgacatg tctggaaagc cgaagagatt 960
tggccgtgct cgcaagagaa ccggaacaca ggtgctgcat ggctgtcgtc agctcgtgtc 1020
gtgagatgtt ggggttaagtc ccgcaacgag cgcaaccctt gtcattagtt gctacgcaag 1080
agcactctaa tgagactgcc ggtgacaaac cggaggaagg tggggatgac gtcaagtcct 1140
catggccctt atgggtaggg cttcacacgt catacaatgg tcgggacaga gggtcgccaa 1200
cccgcgaggg ggagccaatc tcagaaaccc gatcgtagtc cggatcgag tctgcaactc 1260
gactgcgtga agtcggaatc gctagtaatc gcggatcaga atgtcgcggt gaatacgttc 1320
ccgggtcttg tacacaccgc ccgtcacacc atgggagtgg gtttcaccag aagtaggtag 1380
octaacgta aggagggcgc ttaccacggt gggaatcatg actgggag 1428

<210> 4

<211> 1382

<212> DNA

<213> Bacterium-partial 16S rDNA of unknown species from Red Sea coral mucus

<400> 4

gaagaccgct ggcggcaggc ctagcacatg caagtcgagc gcaccttcgg gtgagcggcg 60
gacgggttag taacgcgtgg gaacgtaccc agatctcgg aatagccact ggaaacgggtg 120
agtaataccg catagccct tcgggggaaa gatttatcgg atttgatcg gccgcgtta 180
gattagatag ttggtgggt aatggcctac caagtctacg atctatagct ggttttagag 240
gatgatcagc aacactggga ctgagacacg gccagactc ctacgggagg cagcagtggg 300
gaatcttgga caatgggcgc aagcctgatc cagccatgcc gcgtgagtga tgaaggccct 360
agggtogtaa agctctttcg ccagggaaga taatgacggt acctggtaaa gaaaccccg 420
ctaactccgt gccagcagcc gcggtaatac ggagggggtt agcgttggtc ggaattactg 480
ggcgtaaagc gcgcgtaggc ggactattaa gtcgagggtg aaatcccggg gctcaacccc 540
ggaactgcct tcgatactgg tagtcttgag gtcgagagag gtgagtggaa ctccgagtgt 600
agaggtgaaa ttcgtagata ttcggaagaa caccagtggc gaaggcggct cactggctcg 660
atactgacgc tgaggtgcga aagtgtgggg agcaaacagg attagatacc ctggtagtcc 720
acaccgtaaa cgatgaatgc cagtcgtcgg gtagcatgct attcgggtgac acacctaacy 780
gattaagcat tccgcctggg gactacggtc gcaagattaa aactcaaagg aattgacggg 840

```

ggcccgacaca agcgggtggag catgtggttt aattcgaagc aacgcgcaga accttaccaa 900
cccttgacat ccogtgaccg ctggagagat ccagtttccc ttccggggcac ggtgacaggt 960
gctgcatggc tgtcgtcagc tcgtgtcgtg agatgttcgg ttaagtccgg caacgagcgc 1020
aactcacacc cttagttgcc agcctttagt gggcactcta ggggaactgc ccgtgataag 1080
cgggaggaag gtgtggatga cgtcaagtcc tcatggccct tacgggttgg gctacacacg 1140
tgctacaatg gcatctacag tgggttaatc caaaaagat gtctcagttc ggattgtcgt 1200
ctgcaactcg acggcatgaa gtcggaatcg ctagtaatcg cgtaacagca tgacgcggtg 1260
aatacgttcc cgggccttgt acacaccgcc cgtcacacca tgggagttgg gtctaccgca 1320
aggccgtgcg ctaacttttg aggcagcggg gccacggtag gctcagcgag tggggtgtcc 1380
cc 1382

```

<210> 5

<211> 1483

<212> DNA

<213> Bacterium- partial 16S rDNA of unknown species from Red Sea coral mucus

<400> 5

```

tgacgtggc ggcgtggttc agactccaag tcgaacggga ctctttaact agcttgcctat 60
ttaaagatga gagtggcgaa cgggtgcgta acacgtaaag aacctacct tatgtctggg 120
atagcccacc gaaaggtgga ttaataccgg atgtgatctc tcttctcatg aagagaatat 180
taaagctggg gaccttcggg cctggcgcat aaggagggct ttgcggccta tcagcttggt 240
ggtgaggtaa cggctcacca aggctaagac gggtagctgg tctgagagga tgatcagcca 300
cactggaact tagacacggt ccagacacct acgggtggca gcagtttcga atctttcaca 360
atgggcgaaa gcctgatgga gcaacgccgc gtgggggatg aaggccttcg ggttgtaaac 420
ccctgtcacc aaggataaaa cgctatctat taatactagg tagcctgatg taacttggag 480
aggaaggagt ggctaactct gtgccagcag ccgcggtaat acagagactc caggcgttat 540
tcggattcac tgggcgtaaa ggggtgcgag gcggtcagat gtgtcaggtg tgaaatactg 600
cagcttaact gtagaactgc acttgaaact atttgactag agtatcgag aggtaagcgg 660
aattccaggt gtagcagtga aatgcgtaga tatctggagg aacaccaatg gcgaaggcag 720
cttactggac gattactgac gctcaggcac gaaagcatgg ggagcgaaag ggattagata 780
cccctgtagt ccatgccgta aacgttggtc actaactggt ggaggattcg accccttcag 840
cggccaagct aacgcgataa gtgaaccgcc tgaggactac ggccgcaagg ctaaaactca 900
aaggaattga cgggggcctg cacaagcggg ggagcatgtg gcttaattcg atgcaacgcg 960
aagaacctta cctaggcttg acatgtggac gaccggggca gagatgccct ttctcttcgg 1020
agcggctaca cagggtgctgc atggctgtcg tcagctcgtg tcgtgagatg tttggttaag 1080
tccagcaacg agcgcaaccc ctgccactag ttgccagcat ttagttgggg actctagtgg 1140
gacaaaactct ctctgagagt gggaagggtg ggacgacgtc aagtcagtat ggcccttacg 1200

```


tctagggctg cacacgtgct acaatgcccg gtacagāggg acgcaatacc gcgaggtgga 1260
gcaaatacctt aaagccgggc ccagttcaga ttggagtctg caactcgact ccatgaagtt 1320
ggaatcgcta gtaatggcgc atcagctacg gcgccgtgaa tacgttccca ggccttgtac 1380
acaccgcccg tcacgttatg gaagcccggg cttgccccga agtatgttag ctaaccctt 1440
gtgggtggcg atgtcctaag gtgaggctgg taactggaac gaa 1483

<210> 6

<211> 730

<212> DNA

<213> Bacterium-16S rDNA of unknown species of bacterium from Negev soil

<400> 6

atgccacgct agcgggatgc tttacacatg caagtogaac ggcagcgca gagagcttgc 60
tctcttggcg gcgagtggcg gacgggtgag taatatatcg gaacgtgcc agtagcggg 120
gataactact cgaaagagtg gctaataccg catacgccct acgggggaaa gggggggatc 180
gcaagacctc tcaactattg agcggccgat atcgattag ctagttggtg gggtaaaggc 240
tcaccaaggc aacgatccgt agctggtttg agaggacgac cagccacact gggactgaga 300
cacggcccag actoctacgg gaggcagcag tggggaattt tggacaatgg gggaaaccct 360
gatccagcca tcccgctgt atgatgaagg ccttcgggtt gtaaagtact tttggcagag 420
aagaaaaggt atccccctaat acgggatact gctgacggta tctgcagaat aagcaccggc 480
taactacgtg ccagcagccg cggtaatacg tagggtgcaa gcgttaatcg gaattactgg 540
gcgtaaagcg tgtgtaggcg gttcggaaag aaagatgtga aatcccaggg ctcaaccttg 600
gaactgcatt ttttaactgcc gagctagagt atgtcagagg ggggtagaat tccacgtgta 660
gcagtgagat gcgtagatat gtggaggaat accgatggcg aagcagcccc ctgggatata 720
ctgacgctca 730

<210> 7

<211> 689

<212> DNA

<213> Bacterium-16S rDNA of unknown species of bacterium from Negev soil

<400> 7

ccccagctct gattcccacc gtggtaagcg cctccttac ggtaggcta cctacttctg 60
gtgaaaccca ctcccatggt gtgacgggcg gtgtgtacaa gaccgggaa cgtattcacc 120
gcgacattct gatccgcgat tactagcgat tccgacttca cgcagtcgag ttgcagactg 180
cgatccggac tacgatcggg tttctgagat tggtccccc tcgcggttg gcgaccctct 240
gtcccagca ttgtatgacg tgtgaagccc taccataag ggccatgagg acttgacgcc 300
atccccacct tcctccggtt tgtcaccggc agtctcatta gaggctctt gcgtagcaac 360
taatgacaag ggttgcgctc gttgcgggac ttaacccaac atctcacgac acgagctgac 420
gacagccatg cagcacctgt gttccggctc tcttgcgagc acggccaaat ctcttcggct 480
ttccagatat gtcaagggtg ggtaagggtt ttcgcgttgc atcgaattaa tccacatcat 540

ccaccgcttg tgcgggtccc cgtcaattcc tttagtittt aatcttgoga cgtactccc 600
caggcgggtca acttcacgcg ttagctgcgc tactaaggcc taacggcccc aacagctagt 660
tgacatcggt tagggcgtgg actaccagg 689

<210> 8

<211> 651

<212> DNA

<213> Bacterium-16S rDNA of unknown species of bacterium from Negev soil

<400> 8

ggatgaacgc tggcggcagg cctaacacat gcaagtcgag cggtagagag aagcttgctt 60
ctcttgagag cggcggacgg gtgagtaatg cctaggaatc tgcctggtag tgggggataa 120
cgttcggaaa cggacgctaa taccgcatac gtcctacggg agaaagcagg ggaccttcgg 180
gccttgogct atcagatgag cctaggtcgg attagctagt tgggtgggta atggctcacc 240
aaggcgacga tccgtaactg gtctgagagg atgatcagtc aacttgaac tgagacacgg 300
tccagactcc tacgggaggc agcagtgagg aatattggac aatgggagaa agcctgatcc 360
agccatgccg cgtgtgtgaa gaaggtcttc ggattgtaaa gcaactttaag ttgggaggaa 420
gggttgtaga ttaatactct gcaattttga cgttaccgac agaataagca cgggctaact 480
ctgtgccagc agccgcggta atacagaggg tgcaagcgtt aatcggaatt actgggcgta 540
aagcgcgcgt aggtgggttag ttaaagttgg gatgtgaaat ccccgggctc aaccctggga 600
actgcattca aaactgactg actagggtat ggtagagggt ggtggaattt c 651

<210> 9

<211> 686

<212> DNA

<213> Bacterium-16S rDNA of unknown species of bacterium from Negev soil

<400> 9

ccccagtcg tgattacacc gtggtaacgg tctcccgaa ggtagacta gctacttctg 60
gtgcaaccca ctcccatggt gtgacgggag gtgtgtacaa ggcccgggaa cgtattcacc 120
gcgacattct gattcgcgat tactagcgat tccgacttca cgcagtcgag ttgcagactg 180
cgatccggac tacgatcggg ttgtgtggat tagctccacc tcgcggttg gcaaccctct 240
gtaccgacca ttgtagcacg tgtgtagccc aggcgtaag ggccatgatg acttgacgtc 300
atccccacct tctccgggtt tgtcaccggc agtctcctta gagtgccac cattacgtgc 360
tggtaaactaa ggacaagggt tgcgctcggt acgggactta acccaacatc tcacgacag 420
agctgacgac agccatgcag cacctgtctc aatgttcccg aaggcaccaa tctatctcta 480
gaaagtcat tggatgtcaa ggccgtgtaa ggttcttcgc gttgcttoga attaaaccac 540
atgtccacc gcttgtgcgg gccccgtca attcatttga gttttaacct tgccggcgta 600
ctccccaggc ggtcaactta atgcgttagc tgcgccacta agagctcaag gctcccaacg 660
gctagttgac atcgtttacg gcgtgg 686

This Page is Inserted by IFW Indexing and Scanning Operations and is not part of the Official Record.

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

☐ **BLACK BORDERS**

☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**

☒ **FADED TEXT OR DRAWING**

☒ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**

☐ **SKEWED/SLANTED IMAGES**

☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**

☐ **GRAY SCALE DOCUMENTS**

☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**

☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**

☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.